



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$y_0, y_1, y_2, \dots, y_n$$

approaches a limit y , when n increases indefinitely, the sequence

$$\phi(y_0), \phi(y_1), \phi(y_2), \dots, \phi(y_n)$$

approaches the limit $\phi(y)$, and hence *the identical sequence*

$$k^{y_0}, k^{y_1}, k^{y_2}, \dots, k^{y_n}$$

*approaches a limit.** This limit is *defined* as k^y .† Hence for all real values of y

$$\phi(y) = k^y \text{ and hence } \phi(x) = \log_k(x).$$

EVANSTON, ILLINOIS, September 21, 1903.

CONVERSE AND OPPOSITE PROPOSITIONS.

By C. M. HIMEL, Baker-Himel School, Knoxville, Tenn.

Wentworth, in the revised edition of his Plane Geometry, page 5, makes the following confusing statements:

"If a direct proposition and its opposite are true, the converse proposition is true; and if a direct proposition and its converse are true, the opposite proposition is true."

"Thus, if it were true that

1. If an animal is a horse, the animal is a quadruped;
2. If an animal is not a horse, the animal is not a quadruped;

it would follow that

3. If an animal is a quadruped, the animal is a horse.

Moreover, if 1 and 3 are true, then 2 would be true."

The statements should read: *Whether a direct proposition is true or not, if the converse is true the opposite is true; if the opposite is true the converse is true.*

Thus, in the above example, if 1 be true, then 3 would follow; if 3 be true, then 2 would be true. Consider the general case:

1. Proposition: If a is b , then c is d .
2. Converse: If c is d , then a is b .
3. Opposite: If a is not b , then c is not d .

We may prove that if either of the last two, irrespective of the first, is true, the other is also true.

First. Suppose you know that the converse is true, and you wish to prove

*This required proof.

†Cf. Stolz, *loc. cit.*, p. 138; Tannery, *Theorie des fonctions d'une variable*, p. 114.

